# HIGHLIGHTS OF HIDDEN SPRINGS WETLANDS mid June to mid July

**Description**: from Dry Creek trailhead on north side of Hidden Springs between Cartwright Road and Seamans Gulch Road, explore edge of central wetland and nearby creek to see a rich diversity of wetland species, including several known only from this locality in the Boise area. <a href="Difficulty">Difficulty</a>: half a mile ramble, on flat ground. Insect repellent recommended. Plants listed in alphabetical order by genus within category. \* indicates native species Plants known only from Hidden Spring wetlands in Boise Front in **ALL CAPS**.

#### **WILDFLOWERS**

\*Fiddleneck (Amsinckia spp.) – species difficult to distinguish

Burdock (Arctium minus) - fall-blooming

**Catchweed, madwort** (*Asperugo procumbens*)

**Chicory** (*Cichorium intybus*)

- \*(?)Wild or prickly-cucumber (Echinocystis lobata) unclear if native or not
- \*Common willow-herb (Epilobium ciliatum) several pubescence forms
- \*Western goldenrod (Euthamia occidentalis [previously in Solidago]) fall-blooming
- \*Cleavers, bedstraw, stickywilly, goose-grass (Galium aparine)
- \*COMMON AVENS (Geum aleppicum) only known locality in Boise

**Field pepperweed** (*Lepidium campestre*)

Common mallow, cheeseweed (Malva neglecta)

Black medick (Medicago lupulina)

**Sweetclover** (*Melilotus officinalis*)

\*Yellow monkeyflower (Erythranthe guttata, previously in Mimulus)

**Catnip** (*Nepeta cataria*)

**Common plantain** (*Plantago major*)

- \*Nuttall's or slender cinquefoil (Potentilla gracilis var. fastigiata)
- \*MACOUN'S BUTTERCUP (Ranunculus macounii)
- \*Blister buttercup, cursed crowfoot (Ranunculus sceleratus var. multifidus)

**Curlydock** (Rumex crispus)

Climbing nightshade, bittersweet (Solanum dulcamara)

**Alsike clover** (*Trifolium hybridum*) – flowers pale pink

**Red clover** (*Trifolium pratense*) – flowers dark pink, subtended by leaves

- \*Broadleaf cattail (Typha latifolia)
- \*Stinging nettle (Urtica dioica) STINGING HAIRS

Moth mullein (Verbascum blattaria)

\*American speedwell or brooklime (*Veronica americana*) – some leaves short-petiolate Water speedwell or pimpernel (*Veronica anagallis-aquatica*) – all leaves sessile

#### **FERN RELATIVES**

<sup>\*</sup>Common or field horsetail (Equisetum arvense) – along creek

# NOXIOUS AND/OR INVASIVE WEEDS (not all in bloom)

Jointed goatgrass (Aegilops cylindrica)

Cheatgrass (*Bromus tectorum*) -- other annual *Bromus* also present, including Japanese brome (*B. japonicus*), meadow brome (*B. commutatus*), and barren brome (*B. sterilis*)

**Canada thistle** (*Cirsium arvense*)

Poison hemlock (Conium maculatum)

Field bindweed, wild morning-glory (Convolvulus arvensis)

Mouse or wall barley (Hordeum murinum)

Purple loosestrife (Lythrum salicaria) – fall-blooming, recently sprayed

**Reed canarygrass** (*Phalaris arundinacea*)

Feral or Cereal Rye (Secale cereale)

## OTHER GRASSES AND GRASSLIKE PLANTS (parrial list)

- \*Woolly sedge (Carex pellita, previously C.. lanuginosa) perigynia hairy
- \*Awl-fruit sedge (Carex stipata)
- \*Sedge (Carex spp.) other species difficult to distinguish

**Orchard grass** (*Dactylis glomerata*)

- \*Common spike-rush (Eleocharis palustris)
- \*Baltic rush (Juncus balticus)
- \*Common toad rush (Juncus bufonius var. bufonius)

Rabbitsfoot grass (Polypogon monspeliensis)

- \*American, three-square, or chairmakers bulrush (Schoenoplectus americanus, previously included in Scirpus)
- \*Panicled or small-fruited bulrush (Scirpus microcarpus)
- \*PALE BULRUSH (Scirpus pallidus)

**Rattail fescue** (*Vulpia/Festuca myuros*)

### SHRUBS, TREES, AND WOODY VINES (mostly not in bloom)

- \*Red-osier dogwood (Cornus stolonifera)
- \*Black hawthorn (Crataegus douglasii)

**Russian-olive** (*Elaeagnus angustifolia*) – invasive non-native

**Black walnut** (Juglans nigra)

- \*Black cottonwood (Populus trichocarpa) other species and/or hybrids possibly present
- \*Golden currant (Ribes aureum)

**Dog rose** (*Rosa canina*) – non-native, invasive in some spots

Obtuse- or Round-leaf dog rose (Rosa obtusifolia) – like dog rose, but leaflets hairy

- \*Interior or Wood's rose (Rosa woodsii subsp. ultramontana) native, with slender prickles
- \*Willows (Salix spp.) several species locally, most are difficult to distinguish
- \*Western poison-ivy (Toxicodendron rydbergii, alternatively Rhus radicans) CAUSES DERMATITIS